

Don't Fight COVID-19 Alone

We now have the tools to fight back against this virus. Be your best advocate and reach out to your health care provider to see which medication is right for you.



Overview

COVID-19 Treatments

- Reduce death and hospitalization in people at high risk.
- Should be given as soon as possible:
 - **Oral Antivirals:**
Should be given within five days after onset of symptoms.
 - **IV Treatments:**
Should be given within seven days after onset of symptoms.

For more information, please visit our website:

Michigan.gov/COVIDTherapy

Or scan the QR code:



Tested Positive for COVID-19 ?

Consider COVID-19
Treatments.





Know Your Options

Your health care provider can prescribe the best medication for you depending on your medical history and the length of time you have had COVID-19 symptoms.

Ask Your Doctor!

Who is considered high risk for hospitalization of COVID-19?

Those who are immunocompromised or not up to date on the COVID-19 vaccine are particularly at risk and those with any of the following conditions:

- Older age (e.g., ≥65 years of age)
- Obesity or being overweight
- Pregnancy
- Diabetes
- Immunosuppressive disease or receiving immunosuppressive treatment
- Cardiovascular disease (including congenital heart disease, coronary artery disease, stroke) or hypertension
- Chronic lung diseases (e.g., COPD, moderate to severe asthma, etc.)

For more information about who is considered at high risk of COVID-19, please visit the CDC website by scanning the QR code:



When Given Oral Medication

Complete the whole course of treatment. Do not stop because you feel better.

Do not share the treatment with another individual.

Discuss current medications with your prescribing health care provider and pharmacist to ensure there is no interaction between existing medications and the new one.

