

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. -- = <100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01					M	F	9	10	11	12	Black	White	Hisp
VEHICLE SAFETY																
88	Of students who rode bicycle during past 12 months, % who never/rarely wore a helmet	Bicycle no helmet 12 months	92* 91.7 89.1-94.3	89 89.3 87.0-91.5	93* 93.2 91.1-95.3	95* 95.0 93.5-96.4	85* 84.7 81.8-87.6	92 92.4 89.7-95.1	91 91.1 87.9-94.2	91 90.6 86.0-95.1	94 93.6 91.4-95.7	91 91.4 88.0-94.7	92 91.5 87.4-95.6	93 93.2 88.7-97.6	92 91.6 88.8-94.5	93 92.7 87.9-97.4
99	% of students who never/rarely wore seat belt when riding in a car driven by someone else	Car passenger no seat belt ever	8* 7.9 6.6-9.1	8* 8.3 7.0-9.7	14 13.8 11.3-16.3	19* 18.5 16.3-20.8	14* 14.1 12.4-15.8	9 9.0 7.2-10.8	6 6.4 4.9-7.9	8 8.4 6.4-10.3	7 7.0 4.8-9.3	7 7.3 5.6-9.1	8 7.6 5.0-10.1	10 10.2 5.6-14.7	7 6.9 5.8-7.9	11 11.1 6.6-15.6
1010	% of students who during the past 30 days rode one or more times in a car or other vehicle driven by someone who had been drinking alcohol	Car passenger with driver DUI 30 days	30 29.6 27.5-31.8	32 32.1 28.6-35.6	34 33.6 30.9-36.3	37 37.1 33.4-40.9	31 30.7 28.7-32.7	30 29.8 27.1-32.5	29 29.3 26.7-31.9	26 26.1 22.4-29.8	30 29.6 26.3-32.9	31 30.6 26.1-35.1	33 32.8 29.7-35.9	35 34.8 27.8-41.8	28 28.2 25.5-30.9	36 35.5 27.3-43.7
1111	% of students who during the past 30 days drove a car or other vehicle one or more times when they had been drinking alcohol	DUI 30 days	11 10.7 9.0-12.4	13 12.9 10.7-15.1	13 13.0 11.1-14.9	17 16.5 13.8-19.2	13 13.3 11.8-14.8	12 11.8 9.7-13.9	9 9.4 7.5-11.3	6 6.1 4.4-7.8	8 8.1 6.3-10.0	13 12.9 8.6-17.1	17 16.9 13.3-20.6	8 8.4 5.5-11.3	11 11.1 9.4-12.8	15 14.5 9.1-19.8
VIOLENCE																
1212	% of students who carried a weapon, such as a gun, knife or club, on one or more of the past 30 days	Weapon 30 days	15 15.2 13.5-17.0	13 12.5 11.0-14.0	16 16.0 12.9-19.1	19 18.9 16.4-21.5	17 17.4 15.5-19.3	22 22.1 18.3-25.8	8 8.1 6.2-10.0	18 17.5 15.2-19.8	14 14.4 11.8-17.1	14 14 10.0-17.9	14 13.6 10.0-17.3	17 17.0 11.1-23.0	14 14.4 12.6-16.2	23 23.2 16.6-29.7
1313	% of students who carried a gun on one or more of the past 30 days	Gun 30 days	4 4.4 3.3-5.5	5 4.5 3.8-5.1	6 5.6 4.2-7.0	7 7.0 5.4-8.5	6 5.7 4.7-6.7	7 7.3 5.2-9.4	1 1.2 0.5-1.8	5 4.7 3.2-6.3	4 3.7 1.9-5.5	4 3.6 1.9-5.2	5 4.5 2.4-6.7	6 6.1 2.3-10.0	4 3.6 2.7-4.5	11 10.5 4.8-16.2
1414	% of students who carried a weapon, such as a gun, knife or club, on school property on one or more of the past 30 days	Weapon on school property 30 days	5 5.1 3.8-6.4	5 4.9 4.1-5.8	8 7.5 6.0-9.1	8 8.2 6.9-9.5	6 6.4 5.4-7.4	7 6.8 4.8-8.8	3 3.4 2.0-4.7	6 6.2 4.3-8.0	5 4.8 2.8-6.8	5 4.6 2.6-6.5	4 3.6 2.1-5.0	8 8.1 3.6-12.6	4 4.1 2.9-5.3	13 13.1 8.3-17.9
1515	% of students who did not go to school on one or more of the past 30 days because they felt unsafe at school or on their way to or from school	Missed school felt unsafe 30 days	6 5.5 4.2-6.7	7 7.2 5.5-8.9	5 5.4 4.3-6.5	5 5.1 4.0-6.3	7 6.6 5.6-7.6	6 5.8 4.4-7.2	5 4.7 3.4-6.0	8 7.8 5.7-9.8	4 4.3 3.1-5.5	4 4.2 2.4-6.0	4 4.4 2.1-6.7	10 9.7 6.2-13.2	4 4.1 3.1-5.2	11 10.5 6.8-14.2
1616	% of students who had been threatened or injured with a weapon on school property one or more times during the past 12 months	Threat/injury with weapon on school property	10 9.7 8.6-10.8	9 9.0 8.1-10.0	9 9.2 8.3-10.1	9 9.2 8.0-10.5	9 8.9 7.8-10.0	13 12.6 11.0-14.3	7 6.5 5.1-7.8	14 13.7 11.1-16.4	8 8.1 5.4-10.9	8 7.9 5.2-10.7	7 6.9 5.1-8.7	11 11.0 7.8-14.2	9 9.0 7.6-10.3	16 16.0 8.8-23.2
1717	% of students who had property, such as car, clothing, or books, stolen or deliberately damaged on school property during past 12 months	Stolen/damaged property at school 12 months	32 31.5 29.9-33.1	N/A	N/A	N/A	N/A	35 34.5 32.3-36.7	28 28.1 25.4-30.7	34 34.0 30.5-37.4	32 31.9 29.4-34.4	29 29.2 26.3-32.1	29 29.4 24.8-34.0	31 30.6 27.9-33.3	31 31.2 29.1-33.3	37 36.8 28.4-45.2
1818	% of students who were in a physical fight one or more times during the past 12 months	Physical fight 12 months	31 30.8 27.8-33.7	34 33.8 30.9-36.6	35 35.1 32.0-38.2	36 36.4 33.8-39.0	33 33.2 31.8-34.6	37 37.0 32.9-41.2	24 24.0 21.3-26.7	36 36.3 32.2-40.5	31 31.0 26.9-35.1	26 25.8 20.5-31.1	28 27.5 21.6-33.4	41 40.7 30.9-50.6	28 28.3 25.6-31.0	44 44.1 37.1-51.1

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. --<100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01					M	F	9	10	11	12	Black	White	Hisp
19 19	% of students who were injured in a physical fight and had to be treated by a doctor/nurse one or more times during the past 12 months	Injury in fight required treatment 12 months	3 3.1 2.5-3.7	4 3.8 3.2-4.3	3 3.3 2.6-4.0	4 3.9 2.7-5.0	4 4.0 3.6-4.4	4 3.5 2.6-4.5	3 2.5 1.7-3.2	4 3.7 2.2-5.2	3 2.7 1.4-4.0	4 3.5 2.3-4.6	2 1.8 1.0-2.5	4 3.7 2.1-5.3	3 2.6 2.1-3.2	7 7.3 3.0-11.7
20 20	% of students who were in a physical fight on school property one or more times during the past 12 months	Physical fight on school property 12 months	12 12.2 10.2-14.2	11 11.4 9.6-13.3	14 13.5 11.8-15.2	15 15.2 13.3-17.2	13 12.5 11.5-13.5	16 15.7 13.3-18.2	8 8.1 5.8-10.3	17 16.7 14.2-19.3	11 11.4 8.5-14.3	11 10.5 8.3-12.6	8 7.6 5.0-10.3	20 19.9 14.0-25.9	10 9.9 8.4-11.4	25 25.0 19.2-30.9
21 21	% of students who were ever hit, slapped or physically hurt on purpose by their boyfriend/girlfriend during the past 12 months	Physical violence with boyfriend/girlfriend 12 months	11 11.4 9.8-13.0	12 11.8 10.5-13.1	13 12.9 11.7-14.0	N/A N/A N/A	10 9.5 8.9-10.1	13 12.6 10.5-14.8	10 10.1 8.1-12.1	10 10.2 6.8-13.5	9 9.1 6.7-11.4	14 13.9 11.1-16.8	12 12.1 8.1-16.1	16 15.6 13.2-18.0	10 10.2 8.7-11.7	18 17.7 10.0-25.3
DEPRESSION & SUICIDE																
23 23	% of students who ever felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	Sad or hopeless two weeks 12 months	30 30.2 27.3-33.1	27 27.3 25.5-29.0	27 27.4 25.2-29.6	N/A N/A N/A	28 28.3 27.0-29.6	24 24.1 20.9-27.3	37 36.5 32.5-40.4	33 33.0 27.2-38.8	30 29.9 26.2-33.6	27 26.9 23.3-30.5	30 29.7 24.7-34.7	31 31.1 26.7-35.6	30 29.7 26.5-32.9	37 37.3 28.4-46.2
24 24	% of students who seriously considered attempting suicide during the past 12 months	Suicide considered 12 months	18 18.1 15.7-20.4	18 18.1 16.7-19.5	20 20.1 18.2-22.0	24 23.8 21.1-26.4	19 19 17.6-20.4	14 14.1 11.7-16.6	22 21.8 18.7-25.0	20 20.0 15.5-24.6	17 17.1 14.1-20.1	18 18.2 15.2-21.2	15 15.4 11.4-19.3	15 14.6 11.2-17.9	18 18.3 15.4-21.3	24 24.4 13.7-35.1
25 25	% of students who made a plan about how they would attempt suicide during the past 12 months	Suicide planned 12 months	14 14.2 12.3-16.0	15 14.8 13.8-15.8	15 15.2 13.9-16.5	19 18.6 17.0-20.3	15 14.8 13.7-15.9	12 12.3 10.4-14.2	16 15.9 13.3-18.4	16 16.0 12.8-19.1	14 13.5 10.4-16.5	13 13.1 9.7-16.4	12 12.4 9.5-15.4	12 12.0 9.0-15.1	14 14.2 12.0-16.5	21 21.0 12.0-29.9
26 26	% of students who actually attempted suicide one or more times during the past 12 months	Suicide attempted 12 months	11 10.5 9.2-11.8	10 10.2 8.7-11.6	8 7.9 6.9-8.8	10 10.4 9.0-11.8	9 8.8 8.0-9.6	8 7.8 6.5-9.2	13 12.9 10.8-15.0	14 14.0 11.2-16.9	9 8.9 6.9-10.9	10 9.9 7.6-12.3	7 7.0 4.2-9.7	9 8.9 7.1-10.7	10 10.1 8.6-11.6	18 17.6 10.1-25.1
27 27	% of students whose attempted suicide resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse during the past 12 months	Suicide injury treated 12 months	3 3.2 2.5-3.9	4 3.5 2.8-4.2	3 2.9 2.1-3.6	3 3.1 2.5-3.7	3 2.6 2.2-3.0	2 2.4 1.7-3.2	4 3.9 2.8-5.1	4 4.1 2.9-5.2	2 1.9 1.2-2.6	3 3.4 1.8-5.0	3 2.9 1.2-4.6	3 3.3 1.3-5.2	3 3.0 2.2-3.8	7 7.0 0.9-13.2
TOBACCO																
28 28	% of students who ever tried cigarette smoking, even one or two puffs	Cigarettes Ever	60 60.2 57.0-63.4	64 63.5 60.0-67.1	72 72.2 69.8-74.6	75 75.0 71.8-78.2	64 63.9 61.8-66.0	61 60.6 57.8-63.5	60 59.6 54.7-64.4	53 53.2 48.1-58.3	58 58.0 52.5-63.6	65 64.8 59.7-69.9	67 66.8 60.7-73.0	66 66.2 62.2-70.2	59 58.9 54.9-62.8	70 69.8 63.4-76.2
29 29	% of students who smoked a whole cigarette for the first time before age 13	Cigarettes before age 13	21 21.3 17.7-24.8	23 23.2 20.6-25.7	27 26.7 23.4-30.0	27 27.2 24.4-30.0	22 22.1 20.3-23.9	22 22.0 18.5-25.5	20 20.3 15.5-25.0	23 23.1 19.8-26.5	20 19.9 16.5-23.3	21 20.9 13.7-28.2	20 20.4 14.8-26.0	16 16.4 12.4-20.4	22 21.8 18.0-25.7	26 26.2 22.1-30.3
30 30	% of students who smoked cigarettes on one or more of the past 30 days	Cigarettes 1+ 30 days	23 22.6 18.3-26.9	26 25.7 22.6-28.8	34 34.1 30.9-37.4	38 38.2 34.4-42.0	29 28.5 26.5-30.5	21 21.4 16.0-26.9	24 23.7 19.7-27.8	18 17.9 14.5-21.3	20 19.9 16.5-23.3	23 23.3 14.3-32.3	31 30.6 21.4-39.7	10 9.9 6.1-13.7	25 24.8 20.6-29.1	32 31.8 27.0-36.7

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. -- <100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01				M	F	9	10	11	12	Black	White	Hisp	
M1 30	% of students who smoked cigarettes on 20 or more of the past 30 days	Cigarettes frequent 20+ days 30 days	11 11.3 6.8-15.8	13 12.7 10.1-15.4	17 17.4 14.4-20.5	20 19.8 16.6-23.0	14 13.8 12.2-15.4	12 11.7 5.9-17.4	11 11.0 7.2-14.8	7 6.8 4.4-9.1	8 8.2 6.0-10.4	12 12.4 4.9-19.9	19 19.1 8.6-29.6	3 3.2 1.1-5.3	13 12.8 7.8-17.7	14 13.7 8.0-19.4
31 31	% of students who smoked 2 or more cigarettes per day on the days they smoked during the past 30 days	Cigarettes 2+ per day 30 days	16 15.9 11.1-20.8	18 17.7 15.1-20.4	23 23 20.1-25.8	27 26.9 23.6-30.2	19 19.4 17.5-21.3	16 15.8 9.7-21.8	16 16.1 11.9-20.4	12 11.6 7.7-15.4	13 13.1 10.3-15.9	17 16.5 7.9-25.1	24 23.7 13.4-34.0	5 4.7 1.9-7.5	18 17.9 12.7-23.1	18 18.0 12.8-23.2
M2 31	% of students who smoked more than 10 cigarettes per day on the days that they smoked during the past 30 days	Cigarettes 10+ per day 30 days	4 4.1 1.2-6.9	3 3.2 2.3-4.2	5 5.1 4.1-6.2	6 6.9 5.5-8.3	4 4.1 3.3-4.9	5 4.7 0.6-8.8	3 3.4 1.5-5.2	2 2.1 0.8-3.5	2 2.2 0.9-3.6	5 4.5 0.2-8.7	8 7.6 1.4-13.9	1 1.2 0.0-2.3	5 4.5 1.3-7.7	5 5.3 0.0-10.8
32 32	% of students who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	Cigarettes bought 30 days	7 6.7 4.2-9.2	9 9.2 7.1-11.3	NC	NC	N/A	8 8.1 5.0-11.3	5 5.3 3.2-7.4	2 2.3 1.5-3.2	3 2.7 1.7-3.7	6 5.9 3.1-8.6	18 18.0 10.2-25.9	4 4.2 1.5-6.9	7 7.3 4.4-10.1	6 6.3 2.9-9.7
M3 32	% of students less than 18 years of age who were current smokers, and purchased cigarettes at a store or gas station during the past 30 days	Cigarettes under 18 bought 30 days	18 18.2 14.4-22.1	25 25.1 20.0-30.1	NC	NC	19 19.1 16.9-21.3	24 23.7 19.0-28.5	14 13.9 9.9-17.8	13 12.7 8.3-17.1	13 13.3 8.8-17.8	23 22.8 14.9-30.6	NC	NC	17 17.1 12.8-21.5	NC
33 33	% of students who smoked cigarettes on school property on one or more of the past 30 days	Cigarettes school property 30 days	9 9.3 5.7-12.9	9 9.1 7.3-11.0	13 12.8 10.4-15.3	17 17.3 14.7-19.9	10 9.9 8.7-11.1	9 9.4 5.1-13.8	9 9.1 5.8-12.5	7 6.6 4.1-9.1	6 5.6 3.6-7.5	11 10.5 4.5-16.5	15 15.3 5.7-25.0	3 3.4 1.3-5.5	10 10.3 6.3-14.3	17 16.5 11.7-21.4
34 34	% of students who ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days	Cigarettes daily Ever	18 18.4 13.4-23.3	20 20.4 17.1-23.7	27 27.0 23.5-30.4	N/A	20 20.0 18.1-21.9	18 18.2 11.9-24.4	19 18.5 14.3-22.7	13 13.1 9.5-16.7	17 16.5 13.1-20.0	20 19.5 11.8-27.2	26 25.8 14.9-36.7	8 7.5 5.4-9.7	21 20.6 15.1-26.0	16 16.2 9.4-23.0
35 35	% of students who were current smokers and have tried to quit smoking during the past 12 months	Cigarettes tried to quit 12 months	58 58.4 55.2-61.6	64 64.0 59.7-68.4	NC	N/A	N/A	55 54.7 48.8-60.7	62 61.8 57.8-65.8	60 60.3 50.7-69.8	60 59.7 52.6-66.8	55 54.6 47.8-61.4	60 60.2 53.0-67.3	NC	58 58.3 54.8-61.7	NC
89 36	% of students who have ever tried chewing tobacco, snuff, or dip	Chew, snuff, dip Ever	17 16.6 14.1-19.2	20 19.6 16.6-22.6	24 23.7 20.7-26.8	N/A	N/A	24 24.1 19.9-28.3	9 8.7 6.8-10.6	13 12.5 9.8-15.2	14 13.5 9.9-17.1	20 20.4 14.4-26.4	21 21.2 14.7-27.8	9 9.4 6.5-12.3	18 18.1 15.0-21.3	10 10.3 4.8-15.8
36 37	% of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	Chewing tobacco 30 days	7 6.5 5.2-7.9	8 7.7 5.9-9.5	8 8.2 6.6-9.8	8 8.4 6.0-10.9	8 8.2 6.7-9.7	10 10.0 7.3-12.6	3 2.9 2.0-3.8	5 5.4 3.5-7.3	5 5.3 3.0-7.6	7 7.4 2.8-11.9	8 7.9 4.6-11.2	3 2.6 0.9-4.4	7 7.1 5.7-8.6	9 8.7 0.9-16.5
37 38	% of students who used chewing tobacco, snuff, or dip on school property on one or more of the past 30 days	Chewing tobacco school property 30	3* 2.7 1.8-3.5	4 4.0 2.7-5.3	4 3.9 2.8-5.0	4 4.3 2.7-5.8	5* 5.0 3.8-6.2	4 4.3 2.8-5.7	1 0.9 0.3-1.5	2 1.5 0.7-2.3	2 1.8 0.6-3.0	3 3.4 0.6-6.2	4 3.6 1.2-6.1	1 0.7 0.0-1.4	3 2.8 1.8-3.7	7 6.5 0.0-13.0
38 39	% of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	Cigars 30 days	14 13.6 11.8-15.4	15 14.9 13.0-16.9	20 19.6 17.8-21.4	N/A	15 15.2 14.0-16.4	19 18.5 15.5-21.5	8 8.3 7.2-9.4	12 11.5 8.9-14.2	10 9.5 7.2-11.9	14 14.3 10.2-18.5	20 19.5 15.0-24.0	11 10.7 7.6-13.7	14 13.6 11.5-15.7	21 21.4 15.3-27.4

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. --<100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01				M	F	9	10	11	12	Black	White	Hisp	
M4 30,31, 36,38	% of students who used any tobacco during the past 30 days	Any tobacco 30 days	27*	30	39*		34*	29	26	21	25	29	36	15	30	34
			27.4	29.9	39.2	N/A	33.9	28.5	26.3	21.3	24.8	29.2	36.1	14.9	29.8	33.7
			23.3-31.5	26.5-33.3	36.6-41.8		31.8-36.0	23.2-33.7	22.2-30.4	18.2-24.4	21.4-28.2	20.2-38.2	26.8-45.4	10.2-19.6	25.7-33.9	29.4-38.1
ALCOHOL																
39 40	% of students who had at least one drink of alcohol on one or more days during their life	Alcohol Ever	76	77	82	82	78	74	78	67	76	78	84	71	77	83
			75.9	77.4	81.7	81.9	78.2	74.2	77.5	67.0	76.2	78.2	84.3	71.3	76.7	82.9
			74.0-77.7	74.2-80.6	79.4-84.1	78.5-85.3	76.5-79.9	71.5-76.9	75.0-80.0	63.3-70.8	73.0-79.5	74.2-82.1	80.4-88.2	66.2-76.4	74.4-79.0	77.4-88.5
40 41	% of students who had their first drink of alcohol, other than a few sips, before age 13	Alcohol before age 13	27	27	32	35	29	31	22	35	27	24	19	29	26	35
			26.9	26.9	32.2	34.9	29.1	31.0	22.4	35.3	27.1	23.7	18.5	28.7	26.0	34.6
			24.7-29.1	24.6-29.2	28.9-35.5	31.7-38.1	27.5-30.7	28.2-33.7	19.2-25.5	31.8-38.7	24.2-30.0	18.7-28.6	14.8-22.1	23.2-34.2	23.3-28.7	28.8-40.3
41 42	% of students who had at least one drink of alcohol on one or more of the past 30 days	Alcohol 30 days	44	46	49	51	47	42	46	37	42	44	55	33	46	48
			44.0	46.2	48.5	50.5	47.1	42.3	45.6	37.2	41.8	44.2	55.3	32.5	46.2	48.3
			41.2-46.7	42.6-49.8	45.4-51.7	46.1-54.8	44.9-49.3	38.7-45.9	42.7-48.5	33.6-40.8	36.6-47.0	37.6-50.8	49.5-61.0	28.9-36.2	43.8-48.5	41.5-55.0
42 43	% of students who had 5 or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	Alcohol binge 30 days	27	29	30	32	30	28	27	21	24	29	38	13	30	34
			27.4	29.3	29.9	32.4	29.9	27.9	26.8	20.8	24.2	29.0	38.3	12.7	30.3	33.8
			24.1-30.7	25.6-33.1	27.0-32.8	27.9-36.9	27.9-31.9	23.1-32.7	24.0-29.7	17.3-24.3	18.4-29.9	21.2-36.8	31.1-45.6	9.5-15.9	27.5-33.1	27.8-39.8
43 44	% of students who had a least one drink of alcohol on school property on one or more of the past 30 days	Alcohol school property 30 days	5	5	6	7	5	5	4	6	3	4	6	5	4	9
			4.6	4.9	5.8	7.2	4.9	4.9	4.3	6.1	2.5	3.6	6.0	5.2	4.3	9.3
			4.0-5.3	3.9-6.0	4.7-6.9	6.1-8.4	4.4-5.4	3.7-6.1	3.5-5.1	4.5-7.6	1.8-3.2	2.2-4.9	3.0-9.1	3.3-7.1	3.5-5.1	4.2-14.4
OTHER DRUGS																
44 45	% of students who used marijuana one or more times during their life	Marijuana use Ever	44	44	46	48	42	47	41	33	43	48	55	51	43	54
			44.1	44.0	46.4	48.1	42.4	46.7	41.2	33.4	43.0	47.6	55.4	50.8	42.5	54.2
			40.1-48.0	40.1-47.9	42.9-50.0	43.1-53.1	40.5-44.3	42.3-51.0	36.4-46.0	27.7-39.2	38.1-47.9	40.5-54.6	47.2-63.6	42.7-58.9	38.2-46.9	47.3-61.1
45 46	% of students who tried marijuana for the first time before age 13	Marijuana before age 13	12	12	12	12	10	15	8	14	9	12	10	15	11	17
			11.5	11.6	12.3	12.2	10.2	15.0	7.8	13.7	9.4	11.7	10.1	14.9	10.5	17.0
			8.9-14.0	9.9-13.3	9.6-14.9	10.2-14.2	9.0-11.4	11.2-18.9	5.4-10.1	10.3-17.1	7.0-11.7	7.2-16.3	6.2-14.1	10.8-19.0	7.6-13.4	11.4-22.6
46 47	% of students who used marijuana one or more times during the past 30 days	Marijuana 30 days	24	24	26	28	24	26	23	20	24	26	27	28	23	34
			24.0	24.3	25.9	28.2	23.9	25.5	22.5	20.0	24.0	25.8	27.1	27.8	23.0	33.5
			20.1-27.8	21.7-26.8	22.4-29.4	24.8-31.6	22.4-25.4	20.3-30.7	19.3-25.6	15.7-24.3	19.4-28.7	17.6-34.0	20.6-33.6	20.7-35.0	18.8-27.3	26.2-40.9
47 48	% of students who used marijuana on school property one or more times during the past 30 days	Marijuana on school property 30 days	7	6	6	9	5	8	6	6	6	7	8	9	6	14
			7.0	5.6	6.0	8.9	5.4	8.4	5.5	6.2	6.1	7.0	8.4	9.0	6.4	14.3
			4.6-9.3	4.2-7.0	4.9-7.2	7.2-10.6	4.7-6.1	5.2-11.7	3.8-7.1	3.5-8.9	3.1-9.0	3.0-11.0	3.7-13.2	4.8-13.3	3.8-8.9	8.2-20.4
48 49	% of students who used any form of cocaine, including powder, crack, or freebase, one or more times during their life	Cocaine Ever	9	8	8	7	9	9	8	8	7	9	10	3	9	14
			8.7	7.8	8.1	7.4	9.4	9.3	7.8	7.8	6.8	9.4	10.2	3.2	9.4	14.4
			5.2-12.2	6.0-9.5	6.6-9.6	5.6-9.1	8.2-10.6	4.6-14.1	5.1-10.5	5.1-10.6	4.4-9.2	2.9-15.8	2.5-18.0	1.0-5.5	5.5-13.3	9.6-19.2

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. -- <100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01					M	F	9	10	11	12	Black	White	Hisp
49 50	% of students who used any form of cocaine, including powder, crack, or freebase, one or more times during the past 30 days	Cocaine 30 days	4 3.9 2.4-5.4	4 3.6 2.7-4.5	3 3.4 2.6-4.1	4 3.5 2.6-4.4	4 4.2 3.5-4.9	5 4.5 2.7-6.2	3 3.2 1.7-4.7	4 4.1 2.2-6.0	3 2.6 1.3-3.8	4 3.7 1.2-6.2	5 4.6 0.7-8.5	2 2.3 0.4-4.1	4 3.9 2.1-5.7	7 7.4 2.2-12.7
50 51	% of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paint or spray to get high one or more times during their life	Inhalants Ever	13 13.4 11.2-15.5	13 12.8 10.7-14.9	16 15.6 13.5-17.7	22 21.6 19.0-24.3	15 14.7 13.0-16.4	13 12.9 10.5-15.4	14 13.6 11.2-16.0	15 15.2 12.0-18.5	12 12.1 9.1-15.1	11 11.2 7.5-15.0	14 13.7 10.0-17.4	6 6.4 3.6-9.2	15 14.5 12.2-16.7	19 18.6 11.8-25.3
51 52	% of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during the past 30 days	Inhalants 30 days	4 4.3 3.4-5.2	4 3.6 2.9-4.3	4 4.2 3.2-5.2	N/A	5 4.7 3.9-5.5	5 4.5 3.3-5.7	4 4.0 2.9-5.1	6 6.4 4.3-8.5	3 3.1 1.6-4.7	4 4.1 2.5-5.7	3 2.6 0.9-4.2	4 3.8 1.9-5.7	4 4.1 3.0-5.2	8 7.9 2.7-13.2
90 53	% of students who have taken barbiturates without a doctor's prescription one or more times during their life	Barbiturates Ever	17 17.2 12.8-21.6	17 16.9 13.5-20.2	NC	N/A	N/A	15 15.3 10.0-20.6	19 19.0 14.6-23.5	14 14.2 10.7-17.7	17 17.0 13.4-20.5	18 17.8 8.4-27.2	20 20.0 12.0-28.0	4 3.5 1.6-5.5	20 19.8 15.5-24.0	19 19.3 13.3-25.3
91 54	% of students who have taken barbiturates without a doctor's prescription during the past 30 days	Barbiturates 30 days	9 9.3 7.2-11.4	10 9.7 7.6-11.8	NC	N/A	N/A	9 8.9 5.7-12.1	10 9.5 7.5-11.5	9 8.8 6.2-11.4	7 7.2 4.2-10.2	10 10.2 5.4-15.0	11 10.6 6.2-15.0	2 2.0 0.6-3.5	10 10.4 8.3-12.5	14 13.7 6.1-21.2
52 55	% of students who used heroin one or more times during their life	Heroin Ever	4 3.5 2.2-4.7	3 3.3 2.7-3.9	4 3.6 2.4-4.8	N/A	3 3.1 2.7-3.5	4 4.3 2.7-5.9	3 2.5 1.3-3.6	5 4.5 2.8-6.1	2 1.6 0.6-2.7	4 3.6 0.4-6.9	4 3.5 1.5-5.5	2 2.4 0.7-4.1	3 3.4 2.0-4.7	8 8.4 1.6-15.3
92 56	% of students who used heroin one or more times during the past 30 days	Heroin 30 days	3 2.6 1.8-3.3	2 2.2 1.7-2.6	2 1.9 1.4-2.4	N/A	N/A	3 3.0 2.2-3.7	2 1.9 0.9-2.9	3 3.3 2.2-4.4	1 1.4 0.4-2.4	3 2.6 1.0-4.1	2 2.1 0.7-3.5	3 2.7 0.7-4.6	2 2.2 1.5-3.0	7 7.2 0.0-14.3
53 57	% of students who used methamphetamines one or more times during their life	Methamphetamines Ever	8 7.8 5.8-9.8	8 8.2 6.9-9.6	9 9.0 7.2-10.9	N/A	10 9.8 8.3-11.3	8 8.4 5.3-11.5	7 7.0 5.4-8.6	8 8.4 6.0-10.8	5 5.3 3.6-7.1	9 8.9 4.2-13.6	8 7.7 4.3-11.0	3 3.0 1.1-4.8	8 8.4 6.3-10.6	13 13.1 5.9-20.3
93 58	% of students who have used club drugs one or more times during their life	Club drugs Ever	11 10.8 7.4-14.3	15 15.0 12.8-17.3	N/A	N/A	N/A	11 11.4 6.6-16.1	10 10.2 7.8-12.6	8 7.9 5.8-10.0	8 7.9 5.2-10.7	12 11.6 4.8-18.4	17 16.6 10.0-23.1	5 5.1 2.5-7.7	12 11.7 7.8-15.6	16 15.5 8.8-22.2
94 59	% of students who have used club drugs one or more times during the past 30 days	Club drugs 30 days	5 5.1 3.2-7.0	8 8.2 6.6-9.8	N/A	N/A	N/A	6 5.9 2.8-8.9	4 4.2 2.9-5.5	6 5.7 3.7-7.8	2 2.4 1.2-3.5	6 5.5 0.8-10.2	6 6.1 2.1-10.1	2 2.3 0.8-3.8	5 5.2 3.0-7.4	12 12.1 5.9-18.3
55 60	% of students who took steroid pills or shots without a doctor's prescription one or more times during their life	Steroids Ever	4 3.7 2.8-4.7	4 4.3 3.6-5.0	4 4.0 2.9-5.2	5 5.4 4.2-6.6	5 5.0 4.5-5.5	5 4.7 3.3-6.1	3 2.6 1.5-3.6	4 3.9 2.1-5.6	3 2.8 1.7-3.9	3 2.7 0.9-4.6	5 4.6 2.7-6.5	3 2.5 0.9-4.1	4 3.5 2.5-4.5	8 7.6 0.0-15.5
95 61	% of students who have taken steroid pills or shots without a doctor's prescription one or more times during the past 30 days	Steroids 30 days	3 2.5 1.8-3.3	3 2.5 1.8-3.1	3 2.5 1.8-3.3	N/A	N/A	3 3.0 2.2-3.9	2 1.9 0.9-2.8	3 2.6 1.1-4.1	1 1.4 0.6-2.1	3 2.5 1.4-3.6	3 2.7 1.5-4.0	4 3.9 0.7-7.1	2 1.9 1.3-2.6	10 9.5 2.0-17.1

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. -- = <100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01					M	F	9	10	11	12	Black	White	Hisp
56 62	% of students who used a needle to inject any illegal drug into their body one or more times during their life	Illegal drug injection Ever	2 2.4 1.7-3.1	2 2.4 1.7-3.1	2 2.3 1.6-3.0	3 2.9 2.1-3.7	2 2.3 1.9-2.7	2 2.4 1.7-3.0	2 2.3 1.2-3.3	3 3.4 1.5-5.2	1 1.4 0.6-2.1	2 1.8 0.7-2.8	2 2.3 0.6-3.9	3 2.9 0.8-4.9	2 2.0 1.4-2.7	7 7.0 1.7-12.3
57 63	% of students who were offered, sold, or given an illegal drug on school property during the past 12 months	Illegal drug exchange on school property 12	31 31.3 28.4-34.3	36* 35.6 32.9-38.2	34 33.7 30.3-37.1	36* 36.2 32.8-39.5	29* 28.5 26.5-30.5	35 34.6 30.8-38.4	28 28.0 24.7-31.3	29 28.7 24.5-32.8	33 33.0 28.6-37.3	27 27.4 21.8-33.0	37 37.1 32.1-42.2	33 32.8 24.1-41.4	31 30.5 27.7-33.3	44 43.5 36.0-51.1
SEXUAL BEHAVIOR																
96 64	% of students whose parents or other adults in their family have ever talked with them about what they expected them to do or not to do when it comes to sex	Family discussion regarding sex Ever	72 71.7 69.6-73.8	69 69.4 67.2-71.6	69 69.1 67.4-70.8	NC	N/A	68 68.2 65.1-71.3	76 75.6 73.7-77.4	68 68.3 64.5-72.1	73 72.6 69.4-75.9	75 74.6 67.9-81.3	72 72.4 69.9-74.9	82 81.5 74.8-88.1	71 70.5 68.7-72.4	59 59.0 48.4-69.5
22 22	% of students who have ever been physically forced to have sexual intercourse when they did not want to	Forced sex Ever	10 9.9 8.4-11.4	10 9.5 8.2-10.8	NC	NC	8 7.7 6.8-8.6	7 7.4 5.9-8.8	12 12.4 9.7-15.1	9 8.8 7.0-10.5	8 7.5 5.6-9.4	13 13.0 10.3-15.6	10 9.7 5.5-13.8	14 13.6 11.0-16.2	9 8.7 7.1-10.3	14 13.9 6.8-21.0
58 65	% of students who had sexual intercourse	Sexual intercourse Ever	44 43.5 38.4-48.6	40 40.3 36.7-44.0	45 44.6 40.7-48.5	49 48.9 43.9-53.9	46 45.6 43.3-47.9	45 45.1 39.1-51.0	42 41.9 36.2-47.6	29 28.8 23.4-34.2	37 37.3 30.8-43.7	52 51.7 46.8-56.6	61 61.2 50.3-72.1	67 66.7 60.3-73.2	39 38.9 33.6-44.2	53 53.4 45.7-61.2
59 66	% of students who had sexual intercourse for the first time before age 13	Sexual intercourse before age 13	7 6.9 4.8-9.1	5 4.9 3.5-6.3	7 6.9 5.5-8.2	8 8.3 7.1-9.5	7 6.6 5.7-7.5	10 9.7 6.4-13.1	4 4.0 2.5-5.4	9 8.6 5.6-11.7	6 6.0 2.7-9.3	8 8.1 3.6-12.5	4 3.8 1.8-5.9	21 21.1 17.6-24.6	4 3.9 2.6-5.3	15 15.0 8.6-21.4
97 67	Of students who ever had sexual intercourse, % whose first partner was 3 or more years older	First sex with someone 3+ yrs older	29 28.6 24.4-32.9	25 25.0 22.5-27.6	26 25.6 22.9-28.2	N/A	N/A	19 19.4 15.0-23.8	39 38.5 32.0-45.0	37 36.9 28.1-45.8	29 29.0 21.5-36.6	25 25.4 19.9-30.8	26 26.0 17.6-34.4	37 37.4 29.1-45.7	25 25.0 21.4-28.6	-
60 68	% of students who had sexual intercourse with four or more people during their lives	Sexual intercourse 4+ partners	14 13.5 10.6-16.4	11 10.5 8.5-12.5	14 13.7 10.5-16.9	16 16.4 14.1-18.7	14 14.2 13.0-15.4	16 15.5 11.0-20.1	11 11.3 9.2-13.5	8 7.5 5.4-9.5	10 9.7 5.8-13.5	17 17.3 12.7-21.9	21 21.1 14.3-27.8	32 31.8 25.6-37.9	10 9.9 7.2-12.6	17 17.2 10.4-24.0
61 69	% of students who had sexual intercourse during-the last 3 months	Sexual intercourse 3 months	31 31.1 26.6-35.7	30 29.9 27.0-32.8	31 31.4 27.8-35.1	34 34.4 30.2-38.7	33 33.4 31.4-35.4	30 29.5 24.8-34.1	33 32.8 27.1-38.4	17 17.4 13.5-21.3	25 25.3 20.2-30.4	36 36.4 30.5-42.3	50 50.1 40.2-59.9	44 44.3 35.1-53.5	29 28.5 23.9-33.2	38 37.8 31.2-44.4
M5 69	Of students who have ever had sexual intercourse, % not having sexual intercourse during the past 3 months	Abstinence 3 months	29 28.5 24.2-32.8	26 25.5 22.6-28.3	30 29.7 25.6-33.8	30 29.7 26.8-32.6	27 26.7 25.0-28.4	35 34.5 29.5-39.4	22 21.9 16.7-27.2	39 39.4 33.4-45.4	32 32.2 26.1-38.2	30 29.9 19.2-40.5	18 17.9 13.9-22.0	34 33.5 22.0-44.9	27 26.8 22.9-30.7	-
62 70	Of students who had sexual intercourse during the past 3 months, % who drank alcohol or used drugs before last sexual intercourse	Alcohol or drugs before last intercourse 3	24 24.2 20.0-28.5	24 23.7 19.4-28.0	24 24.4 19.7-29.2	33 32.8 28.4-37.2	26 25.6 23.9-27.3	28 27.9 23.5-32.4	21 20.6 15.6-25.6	26 26.3 16.7-35.9	24 23.7 16.1-31.3	22 22.4 15.9-28.9	24 23.5 17.7-29.3	20 19.8 13.9-25.8	25 24.6 19.2-30.0	-
63 71	Of students who had sexual intercourse during the past 3 months, % who used a condom during last sexual intercourse	Condom last intercourse 3 months	63 62.5 55.6-69.3	61 61.0 58.2-63.9	59 58.9 55.9-61.9	58 58.2 53.1-63.3	58 57.9 55.7-60.1	69 69.0 60.9-77.1	57 56.9 49.8-63.9	74 74.3 66.5-82.2	60 60.4 53.7-67.1	64 63.7 53.5-73.9	59 58.9 47.7-70.1	72 71.7 63.7-79.8	61 60.6 52.3-68.9	-

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. -- = <100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01					M	F	9	10	11	12	Black	White	Hisp
64 72	Of students who had sexual intercourse during the past 3 months, % who used birth control pills during last sexual intercourse	Birth control pills last intercourse 3 months	17	22	20	22	18	14	20	9	16	18	21	7	20	-
			17.1	21.7	20.4	22.3	18.2	13.5	20.0	8.7	16.0	17.5	20.7	6.5	20.4	-
			14.0-20.2	17.2-26.2	16.4-24.4	18.1-26.6	16.5-19.9	9.3-17.8	16.8-23.3	3.9-13.6	9.1-23.0	12.0-23.1	14.9-26.6	2.9-10.1	16.8-24.0	-
M6	% of students who have never had sex, haven't had sex in the last three months, or used a condom the last time they had sex	Responsible sexual behavior	89	89	87	86	86	91	86	96	90	87	80	88	89	85
			88.5	88.6	87.3	85.8	86.1	91.1	86.1	95.6	90.3	87.0	79.5	87.8	88.9	85.0
			85.2-91.8	87.1-90.1	85.7-88.9	83.4-88.2	85.0-87.2	87.8-94.3	82.5-89.7	94.1-97.1	87.5-93.0	82.2-91.9	71.5-87.6	82.5-93.1	85.2-92.6	78.3-91.6
65 73	% of students who had been pregnant or gotten someone pregnant one or more times	Pregnancy ever	5	3	5	6	5	3	5	3	3	6	7	11	3	7
			4.5	2.9	4.9	5.8	4.7	3.4	5.4	2.6	3.1	6.0	6.7	11.2	3.0	7.2
			2.8-6.1	2.1-3.8	3.7-6.2	4.9-6.8	4.2-5.2	2.0-4.9	3.3-7.5	1.0-4.2	1.6-4.7	3.0-9.1	3.1-10.2	6.3-16.1	1.5-4.5	3.8-10.5
87 95	% of students who had ever been taught about AIDS or HIV infection in school	HIV/AIDS instruction in school Ever	89	89	86	88	89	88	90	83	92	91	90	84	90	81
			88.6	88.6	86.1	87.9	89.0	87.7	89.7	83.0	92.1	91.2	90.1	84.0	90.1	80.9
			86.6-90.7	86.9-90.3	82.7-89.5	85.5-90.4	87.7-90.3	85.4-90.1	87.3-92.1	79.5-86.6	90.1-94.1	86.4-96.1	87.5-92.7	79.8-88.3	88.3-91.8	74.6-87.2
WEIGHT & NUTRITION																
M7 6, 7	% of students who are at risk for becoming overweight, whose Body Mass Index is >85% and < 95%	Overweight risk	15	13	15		14	15	15	14	16	17	13	21	14	16
			15.0	13.3	14.7	N/A	13.6	15.2	14.7	13.8	16.1	16.9	13.1	20.9	13.7	16.3
			13.6-16.4	12.0-14.6	13.2-16.3		12.8-14.4	13.2-17.3	12.4-17.1	11.2-16.3	14.4-17.8	13.4-20.5	10.5-15.6	17.4-24.3	12.3-15.1	12.9-19.8
M8 6, 7	% of students who are overweight, whose Body Mass Index is > = 95%	Overweight	12	11	10		11	16	9	13	13	11	12	17	11	17
			12.4	10.7	9.9	N/A	10.5	15.7	8.9	12.8	12.8	11.4	11.9	17.3	11.4	16.8
			10.7-14.1	9.3-12.0	8.6-11.2		9.5-11.5	13.7-17.7	6.8-10.9	9.8-15.8	9.9-15.6	9.6-13.3	8.9-15.0	12.3-22.3	10.1-12.7	9.2-24.3
66 74	% of students who described themselves as slightly or very overweight	Overweight perception	31	31	32	30	29	26	37	29	32	33	33	27	32	34
			31.4	30.7	31.7	30.4	29.2	25.8	37.3	28.6	31.6	32.7	33.2	27.2	32.1	33.5
			29.6-33.2	28.8-32.7	29.5-33.9	28.1-32.8	28.0-30.4	23.5-28.0	34.9-39.7	26.0-31.1	28.5-34.7	29.1-36.4	29.3-37.0	22.5-32.0	30.1-34.1	27.3-39.6
67 75	% of students were trying to lose weight	Trying to lose weight	46	46	45	45	46	31	63	45	49	44	48	40	48	43
			46.4	45.8	44.5	44.8	46.0	30.5	62.7	44.8	49.1	43.9	47.9	39.7	47.8	43.3
			43.9-48.9	43.5-48.1	42.4-46.6	42.6-46.9	44.4-47.6	28.3-32.7	59.0-66.5	39.5-50.1	46.4-51.8	40.5-47.3	43.8-52.1	36.2-43.2	45.1-50.5	34.0-52.7
68 76	% of students who exercised to lose weight or keep from gaining weight during the past 30 days	Exercise 30 days	62	61	60	56	60	51	72	64	61	58	62	54	63	62
			61.6	61.4	60.2	55.7	59.9	51.1	72.2	64.2	61.0	58.3	62.4	54.4	62.7	61.9
			59.2-63.9	59.6-63.2	58.1-62.3	52.8-58.7	58.4-61.4	47.5-54.6	68.0-76.4	61.0-67.5	58.3-63.6	54.7-62.0	56.1-68.7	50.3-58.4	60.3-65.1	53.1-70.7
69 77	% of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight during the past 30 days	Diet 30 days	41	43	43		44	26	56	40	42	39	45	30	43	44
			41.2	42.6	42.8	N/A	43.8	26.4	56.3	39.8	41.8	38.6	44.7	30.0	43.3	44.2
			38.8-43.7	40.5-44.8	40.5-45.0		42.0-45.6	24.0-28.8	52.9-59.7	36.6-42.9	39.0-44.6	33.0-44.3	39.9-49.6	24.2-35.8	40.9-45.8	33.9-54.4
70 78	% of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight during the past 30 days	Fast 30 days	13	14	13		14	9	17	15	12	11	12	15	12	17
			12.6	14.3	13.1	N/A	13.5	8.5	16.8	15.1	11.7	11.3	11.5	15.3	11.9	16.9
			10.8-14.4	12.7-15.9	11.6-14.6		12.4-14.6	7.1-10.0	13.7-19.8	11.2-18.9	10.2-13.3	7.8-14.9	8.5-14.6	13.3-17.2	9.9-14.0	10.5-23.3
71 79	% of students who took diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days	Diet pills 30 days	9	10	8		9	7	11	8	8	9	12	8	9	7
			9.0	9.7	8.1	N/A	9.2	6.6	11.4	7.7	7.7	9.2	11.5	8.4	9.0	7.0
			7.5-10.5	8.5-11.0	7.2-9.1		8.2-10.2	5.5-7.7	9.1-13.7	5.9-9.6	6.1-9.3	6.5-11.9	8.5-14.6	6.4-10.5	7.1-11.0	3.8-10.3

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. -- = <100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01					M	F	9	10	11	12	Black	White	Hisp
72 80	% of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days	Vomit/laxatives 30 days	7 6.6 5.4-7.8	8* 7.6 6.5-8.8	6 5.7 4.5-6.8	6 6.1 5.1-7.2	5* 5.4 4.8-6.0	5 5.3 4.0-6.5	8 7.9 6.3-9.6	8 7.9 5.2-10.7	5 5.1 2.8-7.4	6 5.9 4.1-7.7	7 7.0 4.5-9.5	7 7.2 5.3-9.0	6 6.3 4.9-7.6	13 13.2 7.9-18.4
73 81	% of students who drank 100% fruit juices one or more times during the past 7 days	Fruit juices 7 days	81 81.0 79.7-82.3	85 84.8 83.4-86.3	83 83.4 82.2-84.5	N/A	83 83.0 81.8-84.2	81 80.7 78.5-83.0	81 81.4 79.4-83.3	80 79.8 75.8-83.7	81 81.4 78.6-84.1	82 82.4 78.9-85.8	81 81.1 77.7-84.6	80 79.5 75.4-83.7	81 81.4 80.0-82.8	73 73.1 68.8-77.3
74 82	% of students ate fruit one or more times during the past 7 days	Fruit 7 days	85 85.1 83.1-87.1	85 85.1 82.9-87.4	83 83.4 81.4-85.5	N/A	84 84.4 83.2-85.6	83 83.2 80.8-85.5	87 87.2 84.9-89.5	86 85.5 82.6-88.4	85 85.2 81.9-88.4	86 85.7 82.3-89.1	84 84.0 78.9-89.1	79 79.0 75.7-82.3	87 86.5 84.4-88.6	75 75.4 66.7-84.1
75 83	% of students who ate green salad one or more times during the past 7 days	Green salad 7 days	68 68.0 65.0-71.0	70 70.4 67.6-73.1	65 64.5 61.3-67.8	N/A	67 67.1 65.2-69.0	63 63.0 59.5-66.4	73 73.2 70.3-76.0	65 65.2 62.1-68.3	68 68.1 64.1-72.1	70 70.2 64.5-75.9	70 69.6 64.0-75.2	52 52.4 47.9-56.9	71 71.3 68.3-74.4	58 57.8 50.5-65.1
76 84	% of students who ate potatoes one or more times during the past 7 days	Potatoes 7 days	72 72.1 69.4-74.8	76 76.1 73.8-78.3	75 75.2 73.1-77.3	N/A	73 72.6 70.9-74.3	72 72.2 69.0-75.3	72 72.0 69.1-74.9	72 71.7 67.4-75.9	71 71.2 66.8-75.6	72 71.7 67.4-76.1	74 74.4 70.5-78.2	61 61.2 55.2-67.2	75 74.8 72.2-77.4	66 66.2 58.9-73.5
77 85	% of students who ate carrots one or more times during the past 7 days	Carrots 7 days	50 49.5 44.9-54.2	54 54.2 51.0-57.5	50 49.5 45.6-53.4	N/A	47 47.2 44.7-49.7	50 50.2 44.8-55.7	49 48.6 43.4-53.8	48 47.7 43.8-51.6	51 51.2 46.7-55.8	47 47.1 35.0-59.2	52 52.4 45.6-59.2	25 25.2 16.3-34.1	54 54.3 51.2-57.4	42 42.1 34.1-50.2
78 86	% of students who ate other vegetables one or more times during the past 7 days	Other vegetables 7 days	82 82.0 79.7-84.3	83 82.7 80.2-85.1	83 82.5 80.4-84.6	N/A	85 85.2 84.1-86.3	79 78.8 76.5-81.0	85 85.2 82.4-88.1	77 76.7 72.8-80.5	84 83.5 80.6-86.3	85 85.1 82.0-88.3	84 84.1 80.0-88.2	72 72.0 65.3-78.7	84 84.4 82.6-86.1	75 75.0 68.4-81.5
M9	% of students who ate 5 or more serving of fruits and vegetables per day during the past 7 days	5+ per day fruit/veggies 7 days	18 18.4 16.3-20.4	21 20.6 18.8-22.4	19 19.0 17.1-21.0	NC	21 21.4 20.1-22.7	19 18.7 16.9-20.4	18 17.8 14.6-21.1	18 18.4 15.8-21.1	19 19.0 15.7-22.3	18 18.3 13.0-23.5	17 17.3 13.4-21.3	18 17.8 13.1-22.5	18 18.2 15.9-20.5	20 19.9 14.2-25.5
79 87	% of students who drank 3 or more glasses of milk per day during the past 7 days	Milk 3+ per day 7 days	16 15.5 13.6-17.4	20 20.4 18.3-22.4	21 20.8 18.5-23.1	N/A	16 16.4 15.1-17.7	19 19.0 16.7-21.3	12 11.8 8.9-14.6	17 16.7 14.3-19.1	17 17.1 13.0-21.2	14 13.9 9.6-18.3	13 13.2 9.9-16.5	5 5.3 3.0-7.6	18 17.5 15.6-19.5	14 14.4 7.0-21.8
98 88	Percentage of students who drank 12 ounces of soda pop one or more times per day during the past seven days	Soda consumption 7 days	40 40.1 37.6-42.6	N/A	N/A	N/A	N/A	45 45.1 42.0-48.3	35 34.7 31.5-37.9	38 38.3 35.1-41.5	43 43.2 38.3-48.0	40 40.1 36.8-43.5	37 37.4 31.5-43.3	33 33.4 29.3-37.5	42 41.5 38.6-44.4	46 45.6 40.0-51.2

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. -- = <100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01					M	F	9	10	11	12	Black	White	Hisp
PHYSICAL ACTIVITY																
80 89	% of students who participated in the recommended amount of vigorous physical activities[1] during the past 7 days	Vigorous PA 7 days	62 62.3 59.5-65.1	65 64.5 61.4-67.7	64 63.5 61.5-65.4	N/A	65 64.6 63.1-66.1	67 67.1 63.7-70.4	58 57.8 54.2-61.3	67 66.7 63.2-70.2	64 63.9 60.0-67.8	57 57.1 51.9-62.3	61 61.3 53.9-68.6	50 49.8 43.1-56.6	65 65.1 62.4-67.9	54 53.6 47.6-59.6
81 90	% of students who participated in the recommended amount of moderate physical activities[2] during the past 7 days	Moderate PA 7 days	26 25.6 23.9-27.3	27 26.9 25.4-28.4	27 26.8 24.5-29.1	N/A	26 25.5 24.3-26.7	27 27.3 24.3-30.2	24 23.9 21.5-26.3	24 24.3 21.6-26.9	29 29.1 25.9-32.3	23 22.6 18.1-27.1	27 26.7 22.9-30.5	19 19.0 11.6-26.4	27 27.3 25.4-29.1	21 21.1 16.2-26.1
M10 80 81	% of students who did not participate in the recommended amount of weekly physical activity[3] during the past 7 days	Not recmnded amount of PA 7 days	34 33.6 31.0-36.2	31 31.2 28.2-34.1	N/A	N/A	31 31.2 29.7-32.7	30 29.7 26.6-32.8	37 37.3 33.8-40.9	30 29.5 25.3-33.7	32 32.3 29.3-35.3	40 39.7 34.9-44.6	33 32.9 27.0-38.7	45 44.8 38.8-50.8	31 30.9 28.3-33.5	42 42.2 36.7-47.7
M11 80 81	% of students who participated in no vigorous or moderate physical activity during the past 7 days.	No physical activity 7 days	11 11.0 9.0-12.9	10 9.6 7.9-11.2	N/A	N/A	10 9.5 8.8-10.2	11 10.5 8.5-12.4	12 11.5 9.2-13.9	9 8.9 6.9-11.0	11 10.6 7.2-14.1	11 10.7 8.5-12.8	14 14.1 9.4-18.8	18 18.3 13.7-23.0	9 9.4 7.6-11.2	16 16.3 9.8-22.9
82 91	% of students who did exercises to strengthen or tone their muscles on 3 or more of the past 7 days	Toning muscles 3+ times 7 days	50 50.4 47.5-53.4	52 52.2 49.5-54.8	54 54.1 51.9-56.2	52 51.9 48.0-55.7	53 53.4 51.7-55.1	56 55.7 51.6-59.8	45 45.1 41.6-48.5	54 54.0 49.5-58.5	53 53.2 49.8-56.5	46 45.6 41.2-50.1	48 47.9 41.1-54.8	44 43.8 37.9-49.6	52 51.7 48.7-54.6	45 45.2 39.0-51.4
83 92	% of students who watched 3 hours or more of TV per day on an average school day	TV 3+ hours	35 34.8 31.1-38.5	31 30.5 27.0-34.0	37 36.6 32.0-41.3	N/A	38 38.3 36.0-40.6	39 39.2 35.0-43.4	31 30.5 26.5-34.4	38 38.1 34.5-41.8	34 34.4 29.5-39.3	37 37.0 27.4-46.7	28 28.4 22.5-34.3	61 61.4 55.5-67.3	30 29.6 27.2-32.0	51 51.2 44.8-57.6
84 93	% of students who attended physical education PE class one or more days during an average school week	PE 1+ days avg. school week	39* 39.4 33.7-45.0	44 44.1 39.3-48.9	35* 34.5 29.7-39.3	37* 37.2 32.5-41.8	52* 51.7 47.1-56.3	48 48.2 41.3-55.2	30 30.2 24.4-36.0	61 61.2 54.5-67.9	33 33.0 25.4-40.6	26 26.0 18.7-33.3	32 32.1 24.7-39.6	32 32.2 24.6-39.7	40 40.1 33.6-46.7	39 39.1 25.4-52.8
M12 93	% of students who attended physical education PE class daily	PE 5 days avg. school week	28 27.5 21.2-33.8	29 29.4 22.9-35.9	27 26.9 20.4-33.3	29 29.2 25.8-32.7	32 32.2 27.3-37.1	34 33.6 25.6-41.6	21 21.4 15.8-27.0	44 43.9 35.9-51.9	23 23.2 14.7-31.7	18 18.3 12.7-23.9	21 20.9 14.4-27.3	24 23.9 18.1-29.7	28 28.1 20.8-35.3	28 27.9 17.7-38.0
85 94	Of students enrolled in physical education PE class, % who exercise or played sports more than 20 minutes during an average physical education class	PE-exercise/sport 20+ minutes avg. PE class	86 86.3 82.4-90.1	83 83.2 77.7-88.6	83 82.7 78.8-86.6	NC	83 83.4 81.3-85.5	88 87.6 83.3-91.8	85 84.9 78.4-91.3	86 85.9 81.0-90.8	88 87.5 83.2-91.7	85 84.6 78.4-90.9	87 87.4 80.5-94.3	80 79.8 73.2-86.3	88 87.6 83.5-91.7	- - -

¹ Exercised or participated in physical activities for at least 20 minutes that made them sweat and breathe hard on 3 or more of the past 7 days

² Participated in physical activities that did not make them sweat or breathe hard for at least 30 minutes on 5 or more of the past 7 days

³ Participated in at least 20 minutes of vigorous physical activity on 3 or more of the past 7 days and did not do at least 30 minutes of moderate physical activity on 5 or more of the past 7 days

2003 Michigan Youth Risk Behavior Survey: Detailed Results by Item

N/A=Not available. NC=No comparison due to word change. -- = <100 responses. Shaded areas indicate statistically significant ($p < .05$) differences between percentages within a category, e.g., gender. A double-lined, shaded box = the only number that is a statistically significant difference from the other shaded numbers in a category.																
CB#	Item	Behavior	MI 03	MI 01	MI 99	MI 97	US 01	Gender		Grades				Race/Ethnicity		
Q #			* indicates statistically significance difference from US 01					M	F	9	10	11	12	Black	White	Hisp
MEDICAL HEALTH																
88 5	% of students who saw a doctor or health care provider for a check-up or physical exam when they were not sick or injured during the past 12 months.	Wellness visit 12 months	65 65.2 63.4-66.9	66 66.0 63.8-68.2	62 61.7 59.5-63.9	N/A	N/A	66 65.7 63.2-68.2	65 65.0 62.9-67.1	66 65.8 63.6-68.1	66 66.3 62.1-70.5	66 65.6 61.7-69.5	64 63.8 59.4-68.2	64 63.7 58.1-69.2	66 66.2 64.0-68.4	52 52.3 44.9-59.7
99 96	Percentage of students who have ever been told by a doctor or nurse that they have diabetes.	Diabetes diagnosis	5 4.6 3.3-5.9	N/A	N/A	N/A	N/A	5 5.2 3.3-7.2	4 3.8 2.5-5.2	5 5.1 2.5-7.8	5 4.5 2.9-6.2	4 3.6 2.3-4.8	4 4.1 1.9-6.3	6 6.4 4.4-8.4	4 3.9 2.6-5.1	6 6.0 0.8-11.3
100 97	Percentage of students who have ever been told by a doctor or nurse that they have asthma.	Asthma Diagnosis	21 21.1 19.2-23.1	N/A	N/A	N/A	N/A	23 23.0 20.5-25.4	19 19.1 17.1-21.2	21 21.3 17.3-25.3	22 21.9 18.7-25.1	21 20.7 18.0-23.5	20 19.9 17.0-22.9	25 25.2 20.5-29.9	21 20.5 18.5-22.5	17 17.0 11.3-22.6
101 98	Of students who have been told they have asthma, the % of students who had an episode of asthma or an asthma attack during the past 12 months.	Asthma attacks	26 26.1 22.1-30.1	N/A	N/A	N/A	N/A	20 20.3 15.8-24.8	33 33.2 26.9-39.5	25 24.5 15.9-33.0	31 30.5 22.8-38.1	25 25.3 19.8-30.8	23 22.7 15.9-29.4	26 26.2 17.7-34.7	26 26.4 21.9-30.9	-
GAMBLING																
102 99	Percentage of students who bet on both sporting events and card games and gambled on the internet during the past 30 days.	Gambling - 30 days	2 2.4 1.7-3.0	N/A	N/A	N/A	N/A	4 3.7 2.7-4.7	1 0.8 0.4-1.2	3 2.6 1.8-3.3	2 2.1 1.2-3.0	2 2.0 0.9-3.1	2 2.2 0.9-3.5	4 3.5 2.0-4.9	2 1.9 1.3-2.6	5 4.7 0.4-9.0