



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Di-Syston® 8 Emulsifiable Systemic Insecticide
 For Use on Asparagus to Control Asparagus Aphids
 and Thrips.
 EPA Reg. No. 264-734
 EPA SLN. No. MI-060002

24(c) Supplemental Label

RESTRICTED USE PESTICIDE

Due to acute oral, dermal, inhalation, eye and skin toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MICHIGAN

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	INSECT	RATE	REMARKS
Asparagus	Asparagus Aphid Thrips	1 pt/A	Apply specified dosage per acre to fern stage of asparagus when aphid or thrips infestation appears. Apply in sufficient water for thorough coverage of foliage; a minimum of 5 gallons of water per acre by aircraft or 20 gallons per acre by ground.

RESTRICTIONS:

- Do not apply more than 2 applications of Di-Syston 8 per acre per year.
- Allow at least 6 months between last application and harvest.
- Do not enter or allow workers to enter into treated areas during the restricted entry intervals (REI). The REI for foliar application to asparagus is 11 days.
- Do not apply within a minimum of 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- For use under this SLN label do not apply this product through any type of irrigation system.
- For aerial application, human flaggers are prohibited. Use mechanical flaggers or global positioning system (GPS) equipment.
- Do not apply to asparagus that is grown for seed.

This product is toxic to bees exposed to direct application. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Applications to blooming asparagus must be timed to coincide with periods of minimum bee activity, between late evening and midnight.

This pesticide is extremely toxic to birds, mammals, fish, and aquatic invertebrates. Di-Syston 8 Emulsifiable Systemic Insecticide should not be used under this SLN label where impact on listed threatened or endangered species is likely. Consult the federal label for additional restrictions and precautions to protect aquatic organisms. Certain uses of disulfoton may be restricted by a U.S. District Court final order.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift is the responsibility of the applicator. Do not allow this product to drift. Do not make aerial or ground applications during temperature inversions. Do not apply when winds are greater than 10 mph and avoid gusty and windless conditions. Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal.

This label expires April 17, 2011

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Di-Syston is a registered trademark of Bayer.