Western creek chubsucker (*Erimyzon claviformis*) **- endangered**

Habitat:

feeding - small rivers and creeks, headwater areas
- moderate to high gradient
- sand and gravel substrate
- clear water

spawning - riffles or lake outlets over gravel shoals

winter refuge - larger creeks



Lake chubsucker (*Erimyzon sucetta*)

Habitat:

feeding
- larger clear streams, rivers, lakes, and impoundments
- cannot tolerate turbid water
- low gradient
- prefers dense vegetation over substrate of sand or silt mixed with organic debris

spawning
- small clear streams with moderate to high gradient
- sand or gravel substrate; no clayey silt



Northern hog sucker (*Hypentelium nigricans*)

Habitat:

feeding
- gravel or rubble substrate
- riffles and adjacent pools of warm shallow streams
- clear water
- doesn't like turbidity or siltation
- avoids profuse amounts of aquatic vegetation

spawning
- riffles
- shallow gravel substrate
- high gradient

winter refuge
- deeper quieter pools



Black buffalo (*Ictiobus niger*)

Habitat:

feeding
- large rivers
- deep fast riffles
- occasionally shallow overflow ponds and sloughs
- varying turbidity over various substrates

spawning
- shallows
- sometimes flooded areas



Spotted sucker (*Minytrema melanops*)

Habitat:

feeding
- clear warm rivers (pools, backwaters) with little current
- abundant vegetation
- soft substrate with organic debris
- intolerant of turbidity

spawning
- riffles



Silver redhorse (*Moxostoma anisurum*)

Habitat:

feeding
- streams, rivers, lakes, and impoundments
- low current
- pollution and turbidity intolerant

spawning
- swift current in rivers, do not spawn in tributaries
- males territorial
- gravel to rubble substrate



Black redhorse (*Moxostoma duquesnei*)

Habitat:

feeding
- gravel substrate
- clear water, intolerant of siltation, turbidity, and low gradients
- medium size streams
- cooler swifter streams and short rocky pools with current

spawning
- gravelly riffles

winter refuge
- deeper holes



Golden redhorse (*Moxostoma erythrurum*)

Habitat:

feeding - warm medium gradient streams and rivers
- clear riffly streams
- medium size streams and rivers
- tolerates some turbidity and silt

spawning - shallow gravelly riffles

winter refuge - larger streams



Shorthead redhorse (*Moxostoma macrolepidotum*)

Habitat:

feeding
- downstream sections of large rivers, lakes, and impoundments
- rocky substrates
- swift water near riffles
- clear to slightly turbid water

spawning
- gravelly riffles in smaller feeder streams



Greater redhorse (*Moxostoma valenciennesi*)

Habitat:

feeding - large clear streams
- clean sand, gravel, or boulder substrate
- intolerant of excessive turbidity and chemical pollutants

spawning - moderately rapid current



Black bullhead (*Ameiurus melas*)

Habitat:

feeding
- turbid water
- silt bottom
- low gradient small to medium streams, pools, and headwaters of large rivers; also in lakes and impoundments
- can tolerate very warm water and very low dissolved oxygen

spawning
- nest in moderate to heavy vegetation or woody debris and under overhanging banks



Yellow bullhead (*Ameiurus natalis*)

Habitat:

feeding - clear flowing water
- heavy vegetation
- low gradient streams, lakes, and impoundments
- tolerant of low oxygen

spawning - nest under a stream bank or near stones or stumps



Brown bullhead (*Ameiurus nebulosus*)

Habitat:

feeding
- larger streams and rivers, lakes and impoundments
- clear cool water with little clayey silt
- moderate amounts of aquatic vegetation
- sand, gravel, or muck substrate
- not tolerant of turbid water
- tolerant of warm water and low oxygen

spawning
- nest in mud or sand substrate among rooted aquatic vegetation usually near a stump, tree, or rock

winter refuge
- in muddy bottoms



Channel catfish (*Ictalurus punctatus*)

Habitat:

feeding - moderately-clear, deeper waters of rivers, lakes, and impoundments
- sand, gravel, or rubble substrate
- low to moderate gradient

spawning - secluded semi-dark areas such as holes, under banks, log jams, or rocks



Stonecat (*Noturus flavus*)

Habitat:

feeding
- consistent low to moderate gradient flowing water
- rocky riffles of larger streams and smaller rivers
- not tolerant of silt
- tolerant of low oxygen and pollution

spawning
- eggs deposited beneath stones
- shallow rocky areas of streams or lakes



Tadpole madtom (*Noturus gyrinus*)

Habitat:

feeding
- vegetative cover in low-moderate current waters
- muddy substrate with extensive vegetation
- clear waters of streams, rivers, and lakes

spawning
- mostly in rivers, sometimes shallows of lakes
- nests in dark cavities (ex: beneath boards, logs, crayfish burrows)



Flathead catfish (*Pylodictis olivaris*)

Habitat:

feeding
- (young) shallow riffles in fast current
- deep pools with a lot of woody cover
- deep riffles
- low gradient and current
- prefer silt-free substrate
- sometimes feed on shallow riffles

spawning
- secluded shelters or dark places
- gravel or silt-free substrate

winter refuge
- muddy holes in deep water



Grass pickerel (*Esox americanus vermiculatus*)

Habitat:

feeding - juveniles: along shore
- adults: in deeper portions of streams, rivers, lakes, and impoundments
- clear water, little current, dense vegetation
- tolerates low oxygen concentrations

spawning - broadcast spawner over submerged vegetation



Northern pike (*Esox lucius*)

Habitat:

feeding - cool to moderately warm streams, rivers, lakes, and impoundments
- vegetation in slow to moderate current

spawning - submerged vegetation with slow current in shallow water



Muskellunge (*Esox masquinongy*)

Habitat:

feeding - warm, heavily vegetated lakes, stumpy weedy bays, and slow heavily vegetated medium to large rivers
- shallow cool water
- tolerant of low oxygen

spawning - clear shallow waters (15-20") in heavily vegetated areas



Central mudminnow (*Umbra limi*)

Habitat:

feeding - undisturbed clear, low-gradient streams or rivers and lakes and impoundments
- organic debris, muck, or peat substrates
- aquatic vegetation

spawning - floodplain areas, on vegetation