Opportunity Matrix for D.J. Jacobetti Home for Veterans

	2016	2017	2018	2019	2020		20	21		2022			
	FY16Q4	FY17Q4	FY18Q4	FY19Q4	FY20Q4	FY21Q1	FY21Q2		FY21Q4	FY22Q1			FY22Q4
Performance Summary													
Quality Quintile Performance	3	5	4	4	3	3	2	2	3	3	3	2	2
Quality Measures													
Percent of residents whose need for help with activities of daily living has increased (LS)	3	5	3	3	3	3	2	2	2	2	2	3	3
Percent of residents who self-report moderate to severe pain (LS)	1	3	5	4	4	4	3	3	2				
Percent of high-risk residents with pressure ulcers (LS)	5	5	5	4	4	4	4	3	3	2	1	2	1
Percent of residents who lose too much weight (LS)	4	4	4	4	4	3	2	2	3	4	3	3	2
Percent of low risk residents who lose control of their bowel or bladder (LS)	2	2	1	2	2	3	2	2	2	2	2	2	2
Percent of residents who have/had a catheter inserted and left in their bladder (LS)	3	4	4	4	2	1	1	1	1	1	1	1	1
Percent of residents with a urinary tract infection (LS)	3	4	5	4	3	3	3	3	3	3	3	3	3
Percent of residents who have depressive symptoms (LS)	5	5	5	5	4	4	3	3	3	3	2	3	3
Percent of residents who were physically restrained (LS)	5	5	1	1	1	1	1	1	1	1	1	1	1
Percent of residents experiencing one or more falls with major injury (LS)	2	5	3	5	5	5	4	4	4	3	3	3	3
Percent of residents assessed and appropriately given seasonal influenza vaccine (LS)	4	5	4	1	3	3	3	3	3	3	3	3	1
Percent of residents assessed and appropriately given Pneumococcal vaccine (LS)	3	1	2	4	4	4	3	3	2	2	3	3	3
Percent of residents who received an antipsychotic medication (LS)	3	3	3	4	4	4	4	4	4	4	5	5	5
Percent of residents who self-report moderate to severe pain (SS)	3	4	4	4	4	3	2	1	1				
Percent of residents with pressure ulcers that are new or worsened (SS)	5	5	5	1	1	1	1	1	1	1	1	1	1
Percent of residents assessed and appropriately given seasonal influenza vaccine (SS)	4	4	2	2	2	2	2	2	2	2	2	2	4
Percent of residents assessed and appropriately given Pneumococcal vaccine (SS)	3	3	2	4	2	2	1	1	1	1	1	2	2
Percent of residents who newly received an antipsychotic medication (SS)	2	1	4	4	1	3	5	5	5	5	4	2	2
Percentage of residents whose ability to move independently worsened (LS)			4	3	2	3	4	4	3	3	2	3	3
Percentage of long-stay residents who received an antianxiety or hypnotic medication (LS)			1	2	2	2	3	3	4	4	4	4	4
Percentage of long-stay residents who have had a fall during their episode of care (LS)	2	5	5	5	5	4	4	4	4	4	5	5	5
Percentage of long-stay residents who are receiving antianxiety/hypnotics but do not have evidence of phychotic or related conditions in the target period (LS)	3	2	2	3	3	2	2	3	3	2	1	1	1
Percentage of long-stay residents who have behavior symptoms that affect others during the target period (LS)	1	2	3	2	4	4	4	5	5	5	5	5	5

Red - 5th Quintile; Orange - 4th; Yellow - 3rd; Green - 2nd; Blue - 1st Quintile

(*) Quality Composite for this quarter was calculated using only Long Stay measures.